

#	HIS Section	Source	Revision	Comments
1.	All sections	Indicator	Indicators classified according to type	Defined as Process, Outcome or Impact
2.	All sections	Indicator	Source of standards and indicators referenced where known	
3.	1.0 Population	Guideline	proGres used as default source of population where available	
4.	1.0 Population	Guideline	Source of population data / methodology of collection to be recorded in HIS when ProGRES not available	
5.	1.0 Population	Standard	Internationally agreed % estimates used to calculate denominators where required	Default settings (i) women 15-49 (20%) and (ii) preg and lact (4%).
6.	2.0 Mortality	Standard	CMR standard changed to 1.5 deaths / 1000 / month	To meet SPHERE criteria
7.	2.0 Mortality	Standard	U5MR standard changed to 3 deaths / 1000 / month	To meet SPHERE criteria
8.	2.0 Mortality	Indicator	MMR to be presented as annualised rates only, due to instability of indicator	Annualised rates to be presented in monthly reports
9.	2.0 Mortality	Guideline and Tool	Classification of death to follow: direct/underlying/primary nomenclature	HIV/AIDS, Malnutrition, Maternal, and Measles-related deaths to classified as Primary Causes by default
10.	2.0 Mortality	Guideline	Added indicator "Investigation of Maternal deaths" indicator	To enter part of HIS M&E tool to be developed
11.	2.0 Mortality	Tool	"Unknown" added as reason for death	
12.	2.0 Mortality	Tool	Two additional blank fields added as reason for death	

#	HIS Section	Source	Revision	Comments
13.	2.0 Mortality and 3.0 Morbidity	Guideline	Proportional mortality/morbidity for "Others" should not >25%. To be investigated and broken out when this occurs.	Enter as part of HIS M&E tool
14.	3.0 Morbidity	Indicator	Deleted indicator Proportion STI reported among Nationals	
15.	3.0 Morbidity	Indicator	Deleted indicator Prevalence of syphilis among Nationals	
16.	3.0 Morbidity	Guideline	Definition of new and revisit to be simplified	For adaptation and use in Emergencies
17.	3.0 Morbidity	Tool	National OPD and Syndromic STI data disaggregate by sex	To meet MoH requirements
18.	4.0 IPD and Referral	Indicator	Proportion Mortality in IPD changed to Case Fatality Rate	Disaggregated by Crude and Under 5
19.	5.0 Laboratory	Indicator	Deleted indicator HIV prevalence in Blood donors	
20.	5.0 Laboratory	Indicator	Added indicator % donated units screened in quality assured manner	UNGASS indicator
21.	5.0 Laboratory	Indicator	Deleted indicator syphilis prevalence among nationals	
22.	5.0 Laboratory	Indicator	Added indicator monitoring of RDT positivity rates for malaria	
23.	6.0 Disease Control	Indicator	Added TB Treatment Success Rate	Calculated using cohort analysis in HIS database
24.	6.0 Disease Control	Tool	National data for TB and Leprosy program disaggregated by age and sex	To meet MoH requirements
25.	7.0 EPI and Vitamin A	Standard	Vaccination coverage changed to 95%	
26.	7.0 EPI and Vitamin A	Indicator	Erroneous notes on disaggregation to be removed from Program Dropout rate	

#	HIS Section	Source	Revision	Comments
27.	7.0 EPI and Vitamin A	Indicator	“Growth Monitoring coverage” indicator changed to “Growth Monitoring utilisation”	
28.	7.0 EPI and Vitamin A	Indicator	National EPI data disaggregated by age (1 to < 5)	To meet MoH requirements
29.	7.0 EPI and Vitamin A	Indicator	National Tetanus Toxoid data disaggregated by category (Preg, Non-Preg, Other)	To meet MoH requirements
30.	8.0 Nutrition	Indicator	‘New Admissions’ added to Indicator table	For both SFP and TFP
31.	8.0 Nutrition	Indicator	‘Referral Rate’ added to Indicator table	For both SFP and TFP
32.	8.0 Nutrition	Standard	Death, default and recovery rate changed to reference SPHERE standards	For both SFP and TFP
33.	8.0 Nutrition	Tool and Guideline	“Non-cured” added to reason for exit from OTP and SFP	
34.	8.0 Nutrition	Tool	“Notes” column added to Moderate and Severe Malnutrition Registers	
35.	8.0 Nutrition	Tool and Guideline	Tools and guidelines for monitoring of CTC added to Nutrition section: <ol style="list-style-type: none"> <li>1. SC register</li> <li>2. OTP register</li> <li>3. Weekly and monthly report forms</li> <li>4. New section 8.3 in xls sheet and database</li> </ol>	Guidelines under development

#	HIS Section	Source	Revision	Comments
36.	8.0 Nutrition	Indicator	New indicators for CTC added: For SC: 1. Mean length of stay 2. % successfully stabilised For OTP: 1. Mean length of stay 2. Average weight gain 3. death/default/referral rates 4. % requiring inpatient care 5. Coverage	Disaggregated by Marasmus and Kwashiorkor
37.	9.0 Reproductive Health	Guideline	Timing of First ANC visit redefined from < 16 weeks to within 1 <sup>st</sup> trimester	
38.	9.0 Reproductive Health	Guideline	Definition of Complete Antenatal care revised to 4 antenatal visits	
39.	9.0 Reproductive Health	Guideline	Health centre redefined as Emergency Obstetric Care facility	For comparability with UN Process Indicators
40.	9.0 Reproductive Health	Indicator	Still Birth Rate to be expressed as an annualised figure.	Annualised rates to be presented in monthly reports
41.	9.0 Reproductive Health	Indicator	Indicator “Incidence of obstetric complication” changed to “% obstetric complications treated in EmOC facilities”	UN Process Indicator 3
42.	9.0 Reproductive Health	Indicator	Added indicator % deliveries due to caesarian section	UN Process Indicator 4
43.	9.0 Reproductive Health	Indicator	Added indicator % deliveries at EmOC facility	UN Process Indicator 5
44.	9.0 Reproductive Health	Indicator	Indicator “Incidence of abortion” changed to “Abortion ratio”	According to WHO definition
45.	9.0 Reproductive Health	Indicator	Prevalence of Syphilis (ANC) grouped with Coverage of Syphilis screening.	

#	HIS Section	Source	Revision	Comments
46.	9.0 Reproductive Health	Indicator	Indicator “Coverage of Syphilis screening” denominator redefined to number of pregnant women	
47.	9.0 Reproductive Health	Tool	Removed recording of Haemoglobin (Hb) at each ANC visit, and replaced with Gestational Age	
48.	9.0 Reproductive Health	Tool	Added Caesarian section to legend of complications in delivery register	
49.	9.0 Reproductive Health	Indicator	Deleted indicator Timing of Weighing of Newborns	
50.	9.0 Reproductive Health	Indicator	Deleted indicator % abortions to U18s	
51.	9.0 Reproductive Health	Indicator	Deleted indicator % ratio ANC contacts : ANC partners treated	
52.	9.0 Reproductive Health	Indicator	Deleted indicator % ANC visits to Nationals	
53.	9.0 Reproductive Health	Indicator	Deleted indicator coverage ANC deworming	
54.	9.0 Reproductive Health	Indicator	Deleted indicator ITN distribution	
55.	9.0 Reproductive Health	Indicator	Deleted indicator % births to Nationals	
56.	9.0 Reproductive Health	Indicator	Deleted indicator % obstetric complications to Nationals	
57.	9.0 Reproductive Health	Indicator	Replace Contraceptive Prevalence Rate (CPR) with Couple Years of Protection (CYP)	Pending approval
58.	9.0 Reproductive Health	Guideline	Refine ‘Rape incidence rate’ as ‘Reported rape incidence rate’.	
59.	9.0 Reproductive Health	Guideline	International definitions of rape and sexual assault to be clarified and emphasized during training	
60.	9.0 Reproductive Health	Indicator	Added indicator for timing of PEP provision < 72 hours	

#	HIS Section	Source	Revision	Comments
61.	9.0 Reproductive Health	Indicator	Added indicator for % rape survivors given STI presumptive treatment < 2 weeks	
62.	9.0 Reproductive Health	Tool	Disaggregated National ANC, delivery and PNC data by age (<18 and >18)	To meet MoH requirements
63.	10.0 HIV/AIDS	Tool	Disaggregated condom distribution by male and female (condom)	To meet MoH requirements
64.	10.0 HIV/AIDS	Tool	Disaggregated National VCT and PMTCT data by age (<18 and >18) and sex	To meet MoH requirements
65.	10.0 HIV/AIDS	Indicator	Change to condom coverage to condom distribution rate	Units and standard changed to 0.5 condoms / person / month
66.	10.0 HIV/AIDS	Indicator	Deleted indicator VCT uptake rate	
67.	10.0 HIV/AIDS	Indicator	Change definition of VCT and PMTCT post-test counsel and result to monitor entirety of service provision	
68.	10.0 HIV/AIDS	Indicator	Deleted indicator PMTCT uptake rate	
69.	10.0 HIV/AIDS	Indicator	Deleted indicator HIV Prevalence among ANC partners	
70.	10.0 HIV/AIDS	Indicator	Deleted indicator Proportion of HIV deliveries to Nationals	
71.	10.0 HIV/AIDS	Indicator	Deleted indicator Proportion who plan to Exclusive breastfeed	
72.	10.0 HIV/AIDS	Indicator	Deleted indicator Proportion who receive HBC	
73.	10.0 HIV/AIDS	Indicator	Deleted indicator Proportion who plan to use modern method of FP	
74.	10.0 HIV/AIDS	Indicator	Added % children born to HIV+ mothers who are started on CTR	

#	HIS Section	Source	Revision	Comments
75.	10.0 HIV/AIDS	Indicator	Added % children born to HIV+ mothers who are HIV+ at 18m	
76.	10.0 HIV/AIDS	Indicator	Added % adults and children with adv HIV infection who are started on ARV	
77.	10.0 HIV/AIDS	Indicator	Added % tested HIV+ who are started on CTR / total identified (VCT and PMTCT)	
78.	10.0 HIV/AIDS	Tool	Added “referred from” column to VCT client register	Options: Self referral, TB clinic, STI clinic, OPD, IPD, Blood Donation, Other
79.	10.0 HIV/AIDS	Tool	Added “referred to” column to VCT results register	Options: TB clinic, CTC, PMTCT clinic, Nutrition centre, IPD, None, Other
80.	10.0 HIV/AIDS	Tool	Removed physical and emotional state from VCT and PMTCT registers	
81.	10.0 HIV/AIDS	Tool	Removed duplicate obstetric history fields from PMTCT results register	
82.	10.0 HIV/AIDS	Tool	Removed marital status from VCT and PMTCT registers	
83.	10.0 HIV/AIDS	Tool	Removed obstetric history, apgar score and weight from PMTCT L&D register	
84.	10.0 HIV/AIDS	Tool	Postnatal feeding and FP indicators reworded from % women who “ <b>Plan</b> to accept” to % women who “ <b>Choose</b> to accept ”	